

ヒステリー盲の精神力動
——心身症への分析的アプローチ(1)——

毛 塚 満 男

心 身 医 学

第30巻 第2号 別刷

1990年2月1日発行

日本心身医学会

Original

Psychoanalytical Study of Hysterical Blindness

—Psychoanalytical approach to psychosomatic disorders (1)

Mitsuo Kezuka*

Abstract

Psychosomatic diseases in the field of ophthalmology were classified into three groups by Kato, such as ocular conversion symptoms, ocular symptoms due to neurotic psychization and ocular vegetative neuroses.

Hysterical blindness is the most representative disease of ocular conversion symptoms, therefore many clinical reports have been published.

However, there were no clinical reports discussed in detail from the viewpoint of psychoanalytic theory.

In this paper the recognition gained from a 21-year' old girl suffering from hysterical blindness, who was given supportive psychotherapy based on psychoanalytical understanding, will be described.

As a result, the author has pointed out the psychopathology which involved Oedipal as well as Pre-Oedipal ambivalence in this case.

Received for publication, November 7, 1988.

* Department of Psychosomatic Medicine, Tokyo Komagome Metropolitan Hospital

Address Mitsuo Kezuka: Honkomagome 3-18-22, Bunkyo-ku, Tokyo, 113 Japan.